

Concrete Quarks: The Beginning of the End

G. Zweig, RLE MIT
September 25, 2014
email: zweig@mit.edu

- QCD - developed in two phases:
 - Discovery of quarks
 - Specification of their interactions
- Arose from two very different traditions
 - Rutherford-Bohr
 - Einstein
- Discovery of radioactivity: Henri Becquerel (1896)



Phosphorescence? Becquerel's photographic plate fogged by exposure to radiation from uranium salts. A metal Maltese Cross placed between the plate and the uranium salts is visible.

- Rutherford at Cambridge (1899): α and β

- Rutherford & Soddy at McGill (1903):
“the spontaneous disintegration of [a] radio-element, whereby a part of the original atom was violently ejected as a radiant particle, and the remainder formed a totally new kind of atom with distinct chemical and physical character.”

Nobel prize in Chemistry (1908), Soddy (1921)

PROCEEDINGS OF
THE ROYAL SOCIETY.

SECTION A.—MATHEMATICAL AND PHYSICAL SCIENCES.

On a Diffuse Reflection of the α -Particles.


By H. GEIGER, Ph.D., John Harling Fellow, and E. MARSDEN, Hatfield
Scholar, University of Manchester.

(Communicated by Prof. E. Rutherford, F.R.S. Received May 19,—Read
June 17, 1909.)

In the following experiments, however, conclusive evidence was found of the existence of a diffuse reflection of the α -particles. A small fraction of the α -particles falling upon a metal plate have their directions changed to such an extent that they emerge again at the side of incidence. To form an

We are indebted to Prof. Rutherford for his kind interest and advice throughout this research.

- Interpretation (Rutherford 1911)
 - Impossible!
 - Marsden (1914): Nuclei contain protons!
 - Bohr (1912, 1914-1916): Stationary states
- Charge separation & Quantization



Rutherford's group at Manchester University, 1912.

Rutherford is seated second row, center.

Back rows: (standing): C. G. Darwin, J. M. Nuttall, J. Chadwick,

2nd row: H. Geiger, E. Rutherford,

Front row: H. G. J. Moseley, E. Marsden.

· The nuclear force (1927)

LII. *The Scattering of α -particles by Helium.* By Prof. Sir E. RUTHERFORD, O.M., P.R.S., Cavendish Professor of Experimental Physics, and J. CHADWICK, Ph.D., F.R.S., Fellow of Gonville and Caius College, Cambridge*.

THE study of the collisions of α -particles with hydrogen nuclei has shown that the force between the α -particle and the H nucleus obeys Coulomb's law for large distances of collision, but that it diverges very markedly from this law at close distances. The experiments of Chadwick and Bieler † showed that for distances of collision less than about 4×10^{-13} cm. the force between the two particles increased much more rapidly with decrease of distance than could be accounted for on an inverse square law of force. For

SUMMARY.

into action. Possible explanations of the origin of these additional forces are discussed, and it is suggested tentatively that they may be due to magnetic fields in the nuclei.

Cavendish Laboratory, Cambridge.

· Heisenberg (1925 for atoms; 1943 & 1944 for the nucleus): Work only with observables!

Lone Ranger Atom Bomb Ring Spintharoscope (1947 - early 1950s)

This ring spintharoscope was known as the Lone Ranger Atom Bomb Ring and advertised as a "seething scientific creation." The Lone Ranger was more closely associated with silver bullets than atomic bombs but that's what it was called. When the red base (which served as a "secret message compartment") was taken off, and after a suitable period of time for dark adaptation, you could look through a small plastic lens at scintillations caused by polonium alpha particles striking a zinc sulfide screen.



Distributed by Kix Cereals (15 cents plus a boxtop), the instructions stated: "You'll see brilliant flashes of light in the inky darkness inside the atom chamber. These frenzied vivid flashes are caused by the released energy of atoms. PERFECTLY SAFE - We guarantee you can wear the KIX Atomic "Bomb" Ring with complete safety. The atomic materials inside the ring are harmless."

The following advertisement was appearing in newspapers in early 1947.

Advertisement

SEE GENUINE ATOMS SPLIT TO SMITHEREENS!

INSIDE THIS KIX ATOMIC "BOMB" RING!

HAS SECRET MESSAGE COMPARTMENT

Only 15¢ Plus Boxtop

How Tommy Thwarted the Enemy Agents
with his KIX Atomic "Bomb" Ring

INFO: THE KIX RING WOULD USE EFFECTS OF ATOMIC ENERGY IN ORDER TO BE HARMFUL TO OUR NEW SECRET FORMULA!

OH, THE ENEMY IS BRAGGING AND SAYING "WELL, WE'VE QUARANTINED THE TOP SECRET ATOMIC FORMULA IN THIS KIX RING TO GENERAL HENRICH!"

BUT MY CLIMB HEADQUARTERS I LEARNED THAT A NOTE CARDED BY ENEMY FORMULA. VERY DANGEROUS!

THEY'RE SEEN PLAN! NO? COME HERE!

THEY'RE SEEN PLAN! NO? COME HERE!

THEY'RE SEEN PLAN! NO? COME HERE!

HERE'S HOW TOMMY WAS ABLE TO THWART THE ENEMY AGENTS -- FORMULA WAS HIDDEN IN SECRET MESSAGE COMPARTMENT INSIDE TOP OF KIX ATOMIC "BOMB" RING

HIDDEN ATOMIC CHAMBER IN TOMBARD

CONCEALED OBSERVATION LENS

It's a SEETHING SCIENTIFIC SENSATION!
Actual Atoms—right in smeltorens inside this ring! And you can see brilliant atomic effects! Take Ring into dark room and wait until your eyes are accustomed to darkness. Slide Tail-Fin off—look in Observation Lens—and you'll see frenzied flashes of light—caused by released energy of atoms splitting like crazy. Secret Message Compartment hidden in Tail Fin . . . Tombarrier's Insignia engraved on top of ring. Slides show terrific lightning-blast explosions. Aluminum Warhead (bomb nose) . . . 4-grooved plastic Tail Fin. Ring can be adjusted to fit any finger. Atomic materials in Atom Chamber are harmless. **Now in Get Ring.** Get a line of that winning corn-daddy beautiful cereal KIX. Best top of KIX box together with only 15¢ and Order Blank at right.

HURRY—MAIL THIS ORDER, BLANK TODAY!
SEND TO KIX, Box 65, New York City, N. Y.
Here's my 15¢ plus KIX boxtop. Back to me at once my KIX Atomic "Bomb" Ring.
NAME.....
ADDRESS.....
CITY..... STATE.....
First names and address plainly. Give your number (if any).
NOTE: If you prefer the new "EE Shook 'Em" Ring" package with order blank at top, use that order blank instead of this.
Offer good in U. S. only, while supplies last.

General Mills

THE PHYSICAL REVIEW

A journal of experimental and theoretical physics established by E. L. Nichols in 1893

SECOND SERIES, VOL. 76, NO. 12

DECEMBER 15, 1949

Are Mesons Elementary Particles?

E. FERMI AND C. N. YANG*

Institute for Nuclear Studies, University of Chicago, Chicago, Illinois

(Received August 24, 1949)

The hypothesis that π -mesons may be composite particles formed by the association of a nucleon with an anti-nucleon is discussed. From an extremely crude discussion of the model it appears that such a meson would have in most respects properties similar to those of the meson of the Yukawa theory.

I. INTRODUCTION

IN recent years several new particles have been discovered which are currently assumed to be "elementary," that is, essentially, structureless. The probability that all such particles should be really elementary becomes less and less as their number increases.

It is by no means certain that nucleons, mesons, electrons, neutrinos are all elementary particles and it could be that at least some of the failures of the present theories may be due to disregarding the possibility that some of them may have a complex structure. Unfortunately, we have no clue to decide whether this is true, much less to find out what particles are simple and what particles are complex. In what follows we will try to work out in some detail a special example more as an illustration of a possible program of the theory of particles, than in the hope that what we suggest may actually correspond to reality.

We propose to discuss the hypothesis that the π -meson may not be elementary, but may be a composite particle formed by the association of a nucleon and an anti-nucleon. The first assumption will be, therefore, that both an anti-proton and an anti-neutron exist, having the same relationship to the proton and the neutron, as the electron to the positron. Although this is an assumption that goes beyond what is known experimentally, we do not view it as a very revolutionary one. We must assume, further, that between a nucleon and an anti-nucleon strong attractive forces exist, capable of binding the two particles together.

* Now at the Institute for Advanced Study, Princeton, New Jersey.

We assume that the π -meson is a pair of nucleon and anti-nucleon bound in this way. Since the mass of the π -meson is much smaller than twice the mass of a nucleon, it is necessary to assume that the binding energy is so great that its mass equivalent is equal to the difference between twice the mass of the nucleon and the mass of the meson.

According to this view the positive meson would be the association of a proton and an anti-neutron and the negative meson would be the association of an anti-proton and a neutron. As a model of a neutral meson one could take either a pair of a neutron and an anti-neutron, or of a proton and an anti-proton.

It would be difficult to set up a not too complicated scheme of forces between a nucleon and an anti-nucleon, without about equally strong forces between two ordinary nucleons. These last forces, however, would be quite different from the ordinary nuclear forces, because they would have much greater energy and much shorter range. The reason why no experimental indication of them has been observed for ordinary nucleons may be explained by the assumption that the forces could be attractive between a nucleon and an anti-nucleon and repulsive between two ordinary nucleons. If this is the case, no bound system of two ordinary nucleons would result out of this particular type of interaction. Because of the short range very little would be noticed of such forces even in scattering phenomena.

Ordinary nuclear forces from the point of view of this theory will be discussed below.

Unfortunately we have not succeeded in working out a satisfactory relativistically invariant theory of nucleons among which such attractive forces act. For this reason all the conclusion that will be presented will be

- M. Gell-Mann & E.P. Rosenbaum, “Elementary Particles,” *Scientific American*, July 1957, 72-86: 19 in number

M. Gell-Mann & A.H. Rosenfeld, “Hyperons and Heavy Mesons,” *Ann. Rev. Nucl. Sci*, 1957, 407-478:

Two kinds of Elementary Particles:

Point particles

Spin 1/2 leptons	
Particle	Mass
e^-	1
μ^-	206.7
ν	0

Spin 1 photon	
Particle	Mass
γ	0

Extended particles (strongly interacting)

Spin 1/2 baryons		
Multiplet	Particle	Mass (m_e)
Ξ	Ξ^0	?
	Ξ^{-1}	2585
Σ	Σ^{-1}	2341
	Σ^+	2325
	Σ^0	2324
Λ	Λ	2182
N	n	1838.6
	p	1836.1

Spin 0 mesons		
Multiplet	Particle	Mass
π	π^+	273.2
	π^{-1}	273.2
	π^0	264.2
K	K^+	966.5
	K^-	966.5
	K_1^0	965
	K_2^0	965

– No resonances mentioned!




FIG. 3. A plot of $\pi^\pm + p$ total cross-sections as a function of energy, using in general

- Caltech:
 - Bob Christy ... Alvin Tollestrup
 - My thesis: A test of time reversal symmetry
 $K^+ \rightarrow \pi^0 + \mu^+ + \nu.$
 - Mexico!
 - Murray?


- Every Thursday at 1:30 PM during 1962-63

- Theoretical physics:
 - Axiomatic field theory (no physics)
 - Theory related to belief (Chew, June 1961):

“I believe the conventional association of fields with strongly interacting particles to be empty. ... field theory..., like an old soldier, is destined not to die but just fade away.”
 - Theory related to experiment:
 - * Classification (no dynamics):
 - Sakata model: Wrong baryon spectrum
 - G(2) & SU(3) were in contention

* Dynamics (no classification): Bootstrap

Fred Zacharisen (1961)



Exchanging a ρ binds two pions into a ρ .

But cannot bootstrap the π !

- Experimental physics:

- More particles discovered since 1957:

- * Point particles: the 4th lepton (ν_μ)

- * Extended particles: the 8th spin 1/2 baryon (Ξ^0), and an 8th spin 0 meson (η)

- * Resonances: 26 meson resonances listed in the RMP, April 1963 (ρ , ω , K^* , \dots)

- One Thursday afternoon:

P.L. Connolly, et al., “Existence and Properties of the ϕ Meson”, *Phys. Rev. Lett.* **10**, 371 (1963):




FIG. 1. Dalitz plot for the reaction $K^- + p \rightarrow \Lambda + K + \bar{K}$. The effective-mass distribution for $K\bar{K}$ and for ΛK^+ are projected on the abscissa and ordinate (see reference 7).

$$\phi \rightarrow K\bar{K}$$




FIG. 4. The $M(\pi^+\pi^-\pi^0)$ distribution from the reaction $K^- + p \rightarrow \Lambda + \pi^+ + \pi^- + \pi^0$ after removing Y_1^* production events (see text).

$$\phi \not\rightarrow \rho + \pi$$

$$\begin{aligned} \frac{\Gamma_{K\bar{K}}}{\Gamma_{\rho\pi}} &\sim \left(\frac{p_{K\bar{K}}}{p_{\rho\pi}} \right)^3, \\ &= 1/4 \text{ (expected),} \\ &\geq 35 \text{ (observed).} \end{aligned}$$

“The observed rate [for $\phi \rightarrow \rho + \pi$] is lower than ... predicted values by one order of magnitude; however the above estimates are uncertain by at least this amount so that this discrepancy need not be disconcerting.”

– Feynman:

– GZ:

- Assume hadrons have constituents a called aces:

$$\begin{aligned} &[N_0, \Lambda_0] \quad \& \quad [\bar{N}_0, \bar{\Lambda}_0] \\ &[(p_0, n_0), \Lambda_0] \quad \& \quad [(\bar{p}_0, \bar{n}_0), \bar{\Lambda}_0] \end{aligned}$$

Meson $\equiv a\bar{a}$ with $\uparrow\downarrow$ (e.g. π or K) and
 $\uparrow\uparrow$ (e.g. ϕ or ρ).

Vector mesons as “deuces”





FIG. 2, CERN report TH-401, January 1964.

- A rule for *decay* (in modern notation):



Meson decay: a is an ace, \bar{a} an antiace.

– Implies $\phi \not\rightarrow \rho + \pi$

- A rule for meson masses:

Mass = Σ *constituent masses* + *energies of interaction*, $|\Delta m| > |\Delta E|$.

– Identical binding energies:

$$m^2(\rho) \approx m^2(\omega) < m^2(K^*) < m^2(\phi).$$

$$750^2 \quad 784^2 \quad 888^2 \quad 1018^2$$

– $E_{\Lambda_0}^{\bar{N}_0} = E_{N_0}^{\bar{\Lambda}_0} \approx \frac{1}{2}(E_{\Lambda_0}^{\bar{\Lambda}_0} + E_{N_0}^{\bar{N}_0}), N_0 = p_0, n_0 :$

$$m^2(\phi) \approx 2m^2(K^*) - m^2(\rho).$$

$$1018^2 \quad 1007^2$$

Two birds with one stone.

- Make baryons from 3 aces aaa , not $aa\bar{a}$ (Sakata).

$$B = \frac{1}{3},$$

$$Q = e[I_z + \frac{B+S}{2}],$$

$$[(p_0, n_0), \Lambda_0] \rightarrow [(\frac{2}{3}, -\frac{1}{3}), -\frac{1}{3}].$$

$$3 \times 3 \times 3 = 1 + 8 + 8 + 10.$$

Would you have believed?

More specifically:

- Hadrons have point constituents
- Aces \leftrightarrow Leptons
- Origin of SU(3) symmetry
- Beyond SU(3) symmetry:
 - Restricted representations, quantum numbers:
 - * Baryons only in 1, 8, 10,
Mesons only in 1, 8, and 9.
 - * There is an \vec{L} and an \vec{S} , with $\vec{J} = \vec{L} + \vec{S}$.
 - * $L = 0$ baryons: $(8, J^P = \frac{1}{2}^+)$ and $(10, \frac{3}{2}^+)$,
 $L = 0$ mesons: $(8, J^{PC} = 0^{-+})$ and $(9, 1^{--})$.
 - * Higher L excitations.
 - * $\vec{L} \cdot \vec{S}$ interactions.
 - * 0^{--} ; 0^{+-} , 1^{-+} , \dots forbidden for any L .

- Mass differences break SU(3) & SU(2) symmetry
 - SU(3): $m(p_0) = m(n_0) < m(\Lambda_0)$,
 - SU(2): $m(p_0) < m(n_0)$.

Baryons as “treys”





FIG. 3

CERN report TH-412, February 1964

- Interactions: *Aces, not hadrons, interact.*
 - Strong interaction couplings: “Zweig’s rule” (what’s allowed!)



Graphical representation of the meson-baryon coupling.

The “little loop” encloses antisymmetrized aces.

The subscript “0” on aces is suppressed.

- Electromagnetic and weak currents:

$$a \rightarrow a + \gamma$$

$$a \rightarrow a' + e^- + \nu$$

(Like the “current-quark” model)

- Hierarchy of meson and baryon mass relations

Assume only two-body forces

– Example, baryon octet ($\uparrow\uparrow\downarrow$):

$$m(n) - m(p) + m(\Xi^-) - m(\Xi^0) = m(\Sigma^-) - m(\Sigma^+).$$

$$7.3 \pm 1.3$$

$$8.3 \pm 0.5$$

$$\{8.14 \pm 0.21\}$$

$$\{8.08 \pm 0.08\}$$

$$\{\Delta = 0.06 \pm 0.22\}$$

Like the “constituent-quark” model, but no potential function assumed

(Not the naive quark model!)

- Many additional mass relations for:
 - Pseudoscalar meson octet ($\uparrow\downarrow$)
 - Baryon decuplet ($\uparrow\uparrow\uparrow$)
 - Orbital excited states ($L = 1, 2, \dots$)
(mass splittings with $\vec{L} \cdot \vec{S}$ coupling)

- Cross multiplet & baryon-meson relations:

$$- m(\Xi^*) - m(\Sigma^*) \approx m(\Xi) - m(\Sigma)$$

145 122

$$- m(N) < m(\Lambda) \Rightarrow m(\rho) < m(K^*)$$

940 < 1115 750 < 890

$$- m^2(K^*) - m^2(\rho) \approx m^2(K) - m^2(\pi)$$

0.22 Gev² 0.22 Gev²

- 80 pages

- Not as easy as it looks:

$$- 26 \Rightarrow 7, \text{ exotics}$$

- What did people think? Were aces real?
 - GZ: Aces had dynamics!
 - Murray Gell-Mann:
 - * “Concrete quark model”
 - * Five years after the deep inelastic scattering experiments at SLAC (partons) “Quarks,” *Acta Physica Austriaca, Suppl. IX*, 733-761 (1972)

“In these lectures I want to speak about at least two interpretations of the concept of quarks for hadrons and, the possible relations between them.

First I want to talk about quarks as ‘constituent quarks’.
These were used especially by G. Zweig (1964) [italics added] who referred to them as aces. ...”

More precise to say:

These were introduced by G. Zweig

“The whole idea is that hadrons act as if they are made up of quarks, but the quarks do not have to be real. ...”

That’s a mischaracterization.

“There is a second use of quarks, as so-called ‘current quarks’ which is quite different from their use as constituent quarks ... If quarks are only fictitious there are certain defects and virtues. The main defect would be that we never experimentally discover real ones and thus will never have a quarkonics industry. The virtue is that then there are no basic constituents for hadrons – hadrons act as if they were made up of quarks but no quarks exist - and, therefore, *there is no reason for a distinction between the quark and bootstrap picture: they can be just two different descriptions of the same system, like wave mechanics and matrix mechanics.*” [italica added]

This was Murray’s vision. Concrete quarks?

– Richard Feynman:

- * Current quarks (or aces)?
- * Concrete quarks (really aces)?
 - “The correct theory should not allow you to say which particles are elementary.”
 - Zweig’s rule? Everything that can possibly happen does,
- * “Have I missed anything Zweig?”

Problems with acceptance:

- Aces violated the spin-statistics theorem
 - Rutherford's atom & Bohr's orbits
 - Wegener's continental drift
- Aces violated current dogma:
 - Nuclear democracy
 - Work with observables.

(Copernicus's view of the solar system)

Would you have believed?

Bayes Theorem:

$$P(A|E) = \frac{1}{1+\lambda},$$

where $\lambda \geq 0$, and

$$\lambda = \frac{P(E|\bar{A}) P(\bar{A})}{P(E|A) P(A)}.$$

Since

$$P(\bar{A}) \approx 1 \text{ and } P(E|A) \approx 1,$$

$$\lambda \approx \frac{P(E|\bar{A})}{P(A)}.$$

Acceptance when $P(E|\bar{A}) \ll P(A)$.

- Einstein tradition: $P(E|\bar{A}) \gg P(A)$:
- Rutherford-Bohr tradition: $P(E|\bar{A}) \ll P(A)$

When did acceptance come?

- Pauling
- Dalitz
- Feynman
- Deep inelastic scattering
- ψ/J

Invention or discovery?

Invention: “a product of the imagination.”

Discovery: “the act of finding or learning something for the first time.”

- Current quarks invented (Einstein tradition)
- Aces discovered (Rutherford-Bohr tradition)

google: zweig CERN interview

Conclusion of CERN report TH-412, February 1964

There are, however, many unanswered questions. Are aces particles? If so, what are their interactions? Do aces bind to form only deuces and treys? What is the particle (or particles) that is responsible for binding the aces? Why must one work with masses for the baryons and mass squares for the mesons? And more generally, why does so simple a model yield such a good approximation to nature?

there is also the outside chance that the model is a closer approximation to nature than we may think, and that fractionally charged aces abound within us.